

$$R(x) = \frac{\alpha_0}{n} + \frac{Q(x)}{P'(x)},$$

where

$$Q(x) = \sum_{k=0}^{n-2} x^k \left\{ \alpha_0 a_{k+1} \frac{n-k-1}{n} + \sum_{j=1}^{n-1-k} \alpha_j a_{k+1+j} \right\}.$$

Now $R(x_i) = \alpha_0/n$ if, and only if, $Q(x_i) = 0$ ($i=1, \dots, n$). But Q is of degree $n-2$, hence $Q \equiv 0$. It follows that the coefficients of P must satisfy the equations

$$(3a) \quad \alpha_0 a_{k+1} \frac{n-k-1}{n} + \sum_{j=1}^{n-1-k} \alpha_j a_{k+1+j} = 0 \quad (k=0, 1, \dots, n-2)$$

together with the condition $I(P) = 0$, which is

$$(3b) \quad \sum_{j=0}^n a_j \alpha_j = 0 \quad (a_n = 1).$$

These equations are identical with those obtainable from (2) by the use of Newton's identities which relate the sums of powers of the zeros of a polynomial with the coefficients. However, such an approach presupposes familiarity with these identities, and obscures the interpolatory origin of the formula. Of course, the existence of P does not guarantee the existence of Q_n since we can say (almost) nothing about the zeros x_i except in special cases.

References

1. Z. Kopal, Numerical Analysis, Wiley, New York, 1955.
2. V. I. Krylov, Approximate Calculation of Integrals, trans. A. H. Stroud, Macmillan, New York, 1962.
3. P. L. Tchebichef, Sur les quadratures, J. de Math., 2, 19 (1874) 19.

A NOTE ON THE ALGEBRAIC CLOSURE OF A FIELD

ROBERT GILMER, Florida State University

In order to prove that a given field F has an algebraic closure, it is sufficient to prove the existence of an extension field K of F such that each nonconstant polynomial f over F splits into linear factors over K , for if such a field K exists, then the set of elements of K algebraic over F is an algebraic closure of F [3, p. 194]. To establish the existence of such a field K , Lang in his text *Algebra* [2] proceeds as follows. He first proves that if E is any field, then there is an extension field E_1 of E such that each nonconstant polynomial f over E has a root in E_1 . Applying this result successively to $E_1, E_2, \dots$, Lang obtains an ascending chain $E_1 \subset E_2 \subset \dots$ of extension fields of E such that for each i , every non-

constant polynomial over E_i has a root in E_{i+1} . Then if $L = \bigcup_{i=1}^{\infty} E_i$, it is easy to show that each nonconstant polynomial over E splits into linear factors over L so that L contains an algebraic closure of E . (See [2] pp. 169–170.) The purpose of this note is to observe that in the above process, the field E_1 already contains an algebraic closure of E . This we do by establishing the following result:

THEOREM. *If K is a subfield of a field L and if each nonconstant polynomial with coefficients in K has a root in L , then each nonconstant polynomial with coefficients in K splits into linear factors in $L[X]$.*

Proof. It suffices to prove that for f a nonconstant irreducible monic polynomial with coefficients in K , f is a product of linear factors in $L[X]$. If K has characteristic 0, we define $p = 1$; otherwise p denotes the characteristic of K . (In Bourbaki's terminology [1, p. 71], p is the characteristic exponent of K .) In a splitting field F of f over K , the roots of f all have the same multiplicity p^e for some nonnegative integer e : $f = \prod_{i=1}^n (X - \alpha_i)^{p^e} = \prod_{i=1}^n (X^{p^e} - \alpha_i^{p^e})$. Further, if $f = X^{np^e} + f_{n-1}X^{(n-1)p^e} + \cdots + f_1X^{p^e} + f_0$, and if $\beta_i = \alpha_i^{p^e}$ for each i , $\{\beta_1, \dots, \beta_n\}$ is the set of roots of the irreducible, separable polynomial $g = x^n + f_{n-1}x^{n-1} + \cdots + f_0$ over K [3, p. 120]. The field $K(\beta_1, \dots, \beta_n)$ is a finite, normal, separable extension of K . It is therefore a simple extension of K : $K(\beta_1, \dots, \beta_n) = K(\gamma)$. If h is the minimal polynomial for γ over K , then $K(\gamma)$ is a splitting field for h over K and there is, by hypothesis, a root θ of h in L . The fields $K(\theta)$ and $K(\gamma)$ are each K -isomorphic to $K[X]/(h)$, and hence are K -isomorphic to each other. It follows that $K(\theta)$ is a splitting field of g over K . Therefore, if $p^e = 1$ —that is, if f is separable over K , then L contains a splitting field of f over K . Hence we have proved our theorem in case the algebraic closure of K in L is separable over K . In particular, our proof is complete if K is a perfect field.

In case K is not perfect, we let K_0 be the subfield of L consisting of those elements which are purely inseparable over K . We show that the field K_0 is perfect. Hence if $s \in K_0$ and if $s^{p^e} \in K$, then by hypothesis, $X^{p^{e+1}} - s^{p^e}$ has a root t in L . Therefore, $t^{p^{e+1}} \in K$, $t \in K_0$, and $(t^p - s)^{p^e} = 0$, implying that $t^p = s$ so that K_0 is perfect [3, p. 124]. Further, if $q(x) = \sum_{i=0}^n q_i x^i$ is any nonconstant polynomial over K_0 , then for some positive integer e , $[q(X)]^{p^e} = \sum q_i^{p^e} X^{ip^e} \in K[X]$ so that $v(X)$ has a root α in L . We have $0 = v(\alpha) = [q(\alpha)]^{p^e}$, implying that $q(\alpha) = 0$. We conclude that K_0 is a perfect field with the property that each nonconstant polynomial over K_0 has a root in the extension field L of K_0 . By our previous proof, each nonconstant polynomial over K_0 splits into linear factors in $L[X]$. This property holds then, in particular, for nonconstant polynomials over K , and this completes the proof of our theorem.

References

1. N. Bourbaki, *Éléments de Mathématique, Algèbre*, Book II, Chap. 5, Hermann, Paris, 1950.
2. S. Lang, *Algebra*, Addison-Wesley, Reading, 1965.
3. B. L. van der Waerden, *Modern Algebra*, vol. I, English ed., Ungar, New York, 1948.