

Introduction to Logical Reasoning

Lecture #16

Creating Proofs by Natural Deduction

Professor David Emmanuel Gray

Carnegie Mellon University in Qatar
Northwestern University in Qatar

The Nine Rules of Inference

1. Modus Ponens (M.P.)

$$\begin{array}{l} 1. \ p \rightarrow q. \\ 2. \ p. \\ \hline \therefore \ q. \end{array}$$

2. Modus Tollens (M.T.)

$$\begin{array}{l} 1. \ p \rightarrow q. \\ 2. \ \sim q. \\ \hline \therefore \ \sim p. \end{array}$$

3. Hypothetical Syllogism (H.S.)

$$\begin{array}{l} 1. \ p \rightarrow q. \\ 2. \ q \rightarrow r. \\ \hline \therefore \ p \rightarrow r. \end{array}$$

4. Disjunctive Syllogism (D.S.)

$$\begin{array}{l} 1. \ p \vee q. \\ 2. \ \sim p. \\ \hline \therefore \ q. \end{array}$$

5. Constructive Dilemma (C.D.)

$$\begin{array}{l} 1. \ (p \rightarrow q) \ \& \ (r \rightarrow s). \\ 2. \ p \vee r. \\ \hline \therefore \ q \vee s. \end{array}$$

6. Absorption (Abs.)

$$\begin{array}{l} 1. \ p \rightarrow q. \\ \hline \therefore \ p \rightarrow (p \ \& \ q). \end{array}$$

7. Simplification (Simp.)

$$\begin{array}{l} 1. \ p \ \& \ q. \\ \hline \therefore \ p. \end{array}$$

8. Conjunction (Conj.)

$$\begin{array}{l} 1. \ p. \\ 2. \ q. \\ \hline \therefore \ p \ \& \ q. \end{array}$$

9. Addition (Add.)

$$\begin{array}{l} 1. \ p. \\ \hline \therefore \ p \vee q. \end{array}$$

Natural Deduction

So far we have just seen formal proofs that only require being able to recognize the rules of inference (i.e., the “patterns”) being applied. Now we can begin to begin to construct longer proofs where the steps are not given to us. To introduce you to this process, today we will look at arguments whose proofs can be done in just *two* steps. While these proofs may be short, they will develop the skills necessary for longer and more complicated proofs.

Argument 1

The following is a valid argument. Use natural deduction to construct that argument's formal proof of validity. This proof can be done in only *two* steps.

1. A.

2. B.

$\therefore (A \vee C) \& B.$

Argument 2

The following is a valid argument. Use natural deduction to construct that argument's formal proof of validity. This proof can be done in only *two* steps.

1. $A \rightarrow B.$

2. $A \vee B.$

3. $C \rightarrow D.$

$\therefore B \vee D.$

Natural Deduction

Of course, the real goal with natural deduction is to be able to take arguments in English, translate them into the language of logic, and then formally prove their validity.

Argument 3

The following is a valid argument in English. (1) Translate it into the language of symbolic logic, using the indicated capital letters to label each simple positive statement involved, (2) put it into its argumentative form, and (3) use natural deduction to construct that argument's formal proof of validity. This proof can be done in only *two* steps.

Layli being present is sufficient for Majnun being happy. Nada being pleased is necessary for Layli being present and Majnun being happy. Therefore, Layli is present only if Nada is pleased. (L, M, N)

Learning Natural Deduction

There are only three ways to learn natural deduction:

1. Practice,
2. Practice, and
3. Practice.

If you do not practice this, then you will not be able to do it. I trust you now understand *modus ponens* and *modus tollens*, so you can follow the implications here.

Next Class...

We will look at arguments that require even longer proofs of validity.
Don't panic.

Also, please do not forget to turn in your response to the Lecture #16
Questionnaire on your way out.