



Contents lists available at ScienceDirect

Journal of Steroid Biochemistry & Molecular Biology

journal homepage: www.elsevier.com/locate/jsbmb



Review

Estrogen in prefrontal cortex blocks stress-induced cognitive impairments in female rats

Eunice Y. Yuen, Jing Wei, Zhen Yan*

Department of Physiology and Biophysics, State University of New York at Buffalo, School of Medicine and Biomedical Sciences, Buffalo, NY 14214, USA

ARTICLE INFO

Article history:

Received 11 May 2015
Received in revised form 20 August 2015
Accepted 24 August 2015
Available online xxx

Keywords:

Stress
Sexual dimorphism
Estrogen
Glutamate receptors
Cognition

ABSTRACT

Animal and human studies have found that males and females show distinct stress responses. Recent studies suggest the contribution of estrogen in the brain to this sexual dimorphism. Repeated stress has been found to impair cognitive behaviors via suppressing glutamatergic transmission and glutamate receptor surface expression in pyramidal neurons of prefrontal cortex (PFC) in male rats. On the contrary, female rats exposed to the same stress paradigms show normal synaptic function and PFC-mediated cognition. The level of aromatase, the enzyme for the biosynthesis of estrogen, is significantly higher in the PFC of females than males. The stress-induced glutamatergic deficits and memory impairment are unmasked by blocking estrogen receptors or aromatase in females, suggesting a protective role of estrogen against the detrimental effects of repeated stress.

© 2015 Elsevier Ltd. All rights reserved.

Contents

1. Sexually dimorphic effects of stress and role of estrogen	00
2. Resilience vs. vulnerability of females to stress-related mental disorders	00
3. Role of estrogen in synaptic regulation and brain diseases	00
4. Sexual dimorphism in developing brain	00
Acknowledgement	00
References	00

1. Sexually dimorphic effects of stress and role of estrogen

Corticosteroid stress hormones serve as a key regulator of cognitive and emotional processes [20,43]. It has been proposed that there is an “inverted U” relationship of stress to cognitive function [22], such that a moderate level of corticosteroid has pro-cognitive effects, while too low or too high corticosteroid levels are detrimental to cognitive processing [27]. Our group has found that stress exerts dual effects on cognition through bi-directional modulation of glutamatergic transmission in prefrontal cortex (PFC), a key target region of stress hormones. In young (~4 weeks old) male rats, acute stress significantly enhances glutamate receptor-mediated synaptic currents and improves working memory [75,76]. Conversely, young male rats exposed to one-week repeated restraint or unpredictable stress show the

diminished PFC glutamatergic transmission and the impaired PFC-mediated cognitive function, temporal order recognition memory (TORM, [77]).

While these findings support the notion that short-term (acute) stressors elicit adaptive and beneficial changes, whereas long-term (chronic) stress results in maladaptive and deleterious effects, this pattern of stress responses appears to apply to only males. In response to one acute stressful event of intermittent tail-shocks, spine density is enhanced in the male hippocampus but reduced in the female hippocampus [62]. When the subchronic stress challenge, which induces cognitive impairment in males [77], is introduced to young female rats, their glutamatergic transmission in PFC and TORM function are unaffected [70, Fig. 1A]. Similar sex differences to chronic stress have also been reported by other groups. For example, in male rats, restraint stress (6 h/day, 21-day) impairs performance on a variety of spatial memory tasks including radial arm maze, object placement, Y-maze, water maze, and a nonspatial, recognition memory test [6,7,17,31]. In contrast, females exposed to the same stress paradigm show

* Corresponding author. Fax: +1 7168292699.
E-mail address: zhenyan@buffalo.edu (Z. Yan).




Fig. 1. Estrogen protects against the detrimental effects of repeated stress on cognition. (A) Bar graphs (mean \pm SEM) showing the discrimination ratio (DR) of temporal order recognition memory (TORM) tasks in control or repeatedly stressed (restraint, 7-day, RS) male or female rats (4-wk-old). (B and C) Bar graphs showing the DR of TORM tasks in control vs. repeatedly stressed females with injections of the ER antagonist ICI182,780 (B, 0.05 mg/kg, s.c.), or males with the injections of the ER agonist estradiol (C, 0.1 mg/kg, s.c.). (D and E) Bar graphs showing the DR of TORM tasks in control vs. repeatedly stressed females with the PFC injection of GFP or ER α shRNA lentivirus (D), or with the injections of aromatase inhibitor formestane (2 mg/kg, s.c.). **: $p < 0.005$, ANOVA (A-E). (F) Quantitative real-time RT-PCR data on the mRNA level of ER α , ER β and aromatase in PFC from male vs. female rats. #: $p < 0.05$, *T*-test. Adapted from Wei J, et al., Mol. Psychiatry, 19: 588–598, 2014.

enhanced cognition and memory in almost all of these tasks [7,12,10,9,8,44,18,31]. These animal studies suggest that females are more resilient to chronic stress than males, at least in terms of the measured cognitive behaviors [16].

Interestingly, when estrogen receptor (ER) function is blocked in female rats, the detrimental effects of repeated stress (2-h restraint, 7-days) on PFC glutamatergic transmission and TORM function are unmasked [70], Fig. 1B). On the other hand, when estradiol is administered in male rats, they become resilient to the same stressor [70], Fig. 1C). It suggests that an estrogen-mediated mechanism makes females less susceptible to the deleterious effects of repeated stress than males.

However, the role of estrogen in stress responses is not without controversy. There are also reports suggesting that estrogen may amplify the stress responses in females. Activating stress systems pharmacologically by FG7142, a benzodiazepine inverse agonist, induces impaired PFC working memory in females during proestrus (high estrogen), but not during estrus (low estrogen), suggesting that estrogen may increase the sensitivity to stress in females [60]. Estrogen replacement in ovariectomized female rats exposed to a behavioral stressor (2-h immobilization for 10 days) also induces the greater dendritic remodeling in PFC neurons projecting to the basolateral nucleus of the amygdala (BLA) [61]. Preclinical studies using fear conditioning and extinction paradigms have found that females with low estrogen levels exhibit

impaired extinction retrieval [45,46], which can be reversed by stimulation of D1 dopamine receptors [58]. It suggests that estrogen might influence PFC-BLA function in part through dopaminergic mechanisms.

Converging evidence supports that females and males exhibit different biochemical, cellular and behavioral effects of stress [62,37,11,40,5]. However, the observed sex differences of stress responses and role of estrogen could be affected by a number of factors, such as animal strains, animal ages, stress paradigms, estrogen regimen, and measured outcomes.

2. Resilience vs. vulnerability of females to stress-related mental disorders

Epidemiological studies indicate that women are more likely to develop stress-associated mental disorders, such as depression and PTSD [71,14]. Therefore, it is easy to assume that females have higher stress susceptibility. However, it is important to note that gender vulnerability in stress responses is different from gender vulnerability in mental disorders. Despite the stress exposure for almost everyone, only a small population develops stress-associated mental disorders, including depression and PTSD. Genetic risk factors carried by the susceptible individuals are likely to play a causal role, while stress may only serve as a trigger to precipitate a variety of emotional and cognitive difficulties.

Genetic factors probably also directly influence the intrinsic sensitivity to stress, which contributes to the pathogenesis of psychiatric diseases [28]. Recent studies suggest that individuals carrying Vall66met allele of the *BDNF* (brain derived trophic factor) gene have the altered vulnerability to stress and antidepressant responses [74]. Epigenetic mechanisms involving chromatin remodeling and gene expression could also influence stress sensitivity [65,67].

There are two peaks of depression prevalence in women: postpartum depression, and perimenopausal depression, both of which are associated with large drops (postpartum) or excessive fluctuations (perimenopausal) of serum estradiol levels. Human PET imaging studies suggest that estrogen decline during perimenopausal age elevates the level of monoamine oxidase A, a neurobiological change that also presents during major depressive episodes [57]. The critical role of estrogen in mood disorders is also supported by animal studies. It has been found that injecting estrogen reverses helplessness in animal depression models and enhances hippocampus plasticity [24,13]. Females are more responsive to the action of antidepressants than males [23]. Corroborating with the role of estrogen in females' stress resilience [70], estrogen also shows anxiolytic and antidepressant effects in animal models, which are dependent upon the utilized regimen of estrogen and interactions with the hypothalamic–pituitary–adrenal axis [68].

3. Role of estrogen in synaptic regulation and brain diseases

One surprising finding is that the stress resistance in females is unchanged by ovariectomy that only terminates the ovarian estrogen, but is blocked by knockdown of ER α only in the PFC region [70, Fig. 1D] or global inhibition of aromatase, the estrogen synthesis enzyme [70, Fig. 1E], implying that CNS-synthesized estrogen [72,32] may determine the sexually dimorphic vulnerability to stress. Consistently, it has been shown that estrogen can be synthesized by aromatase localized in neurons from endogenous cholesterol [26]. Moreover, ischemic neuroprotection in females has been attributed to the local, nongonadal estrogen, which may be aromatized from precursor androgens [39]. The expression of aromatase is significantly higher in PFC neurons of female than male rats [70, Fig. 1F], suggesting that pre-pubertal female PFC has an endogenous capacity to generate estrogen that provides protection against repeated stress.

The protective effect of estrogen against stress is in line with the critical role of estrogen in neurogenesis [52], synaptic spine growth [24] and memory consolidation [59]. Depriving estrogen in rats induces postpartum–depression symptoms, and altered expression of genes involved in learning and memory [66]. It has been found that estrogen protects females from several neurological diseases, including seizure [56], ischemia [54] and Alzheimer's disease [36]. Studies suggest that estrogen receptor (ER α), which is expressed at synapses of hippocampus [1] and PFC [69], influences synaptic function through a non-genomic mechanism [41]. For example, estrogen potentiates the activation of NMDA receptors and voltage-gated calcium channels, which results in the increased calcium influx and new excitatory synapses in hippocampus [73,42]. Intracellularly, estrogen rapidly activates synaptic Akt that facilitates local synthesis of PSD-95, a scaffolding protein required for spinogenesis [2]. Moreover, estrogen activates RhoA–ROCK–LIMK–cofilin signalling pathway that controls actin polymerization involved in spine growth [64,33]. At circuitry level, estrogen suppresses inhibitory GABAergic interneuron by downregulating *BDNF* synthesis, which indirectly increases excitatory synaptic transmission of pyramidal neurons in hippocampus [48].

Recent studies have uncovered new functions of brain-derived estrogen (17 α estradiol). This neurosteroid can be produced in

hippocampus [30]. It is proposed that estrogen exerts its action in various brain regions through a combination of genomic and nongenomic mechanisms. Unlike the sustained effect of ovarian estrogen, injection of 17 α estradiol in ovariectomized animals rapidly induces hippocampal spine formation without increasing the expression of synaptic proteins [64]. Electrophysiological data suggests that 17 α estradiol exerts an immediate effect on glutamatergic transmission by enhancing presynaptic glutamate release [63].

4. Sexual dimorphism in developing brain

Evaluation of the distribution of ER α and ER β has revealed their presence in diverse brain regions (cerebral cortex, basal forebrain, amygdala, etc) through early postnatal periods (p3–p14), supports a potential role for estrogens in neural development [53]. Our study shows that estrogen protects prepubertal females from the detrimental effects of repeated stress [70], suggesting that estrogen plays a role in developing female brains, rather than being a typical sex hormone for reproduction. In agreement with this, mounting studies pinpoint the sexual dimorphism in brain development. For instance, comparing to neurogenesis in females, males have a higher proliferation rate and a better survival rate of de novo neurons [78]. Imaging study indicates that white matter development is strongly influenced by hormonal environment of estrogen during early postnatal period, but is minimally affected later in life [34]. Significantly decreased levels of ER β and aromatase (the enzyme converting testosterone to estrogen) have been found in PFC of human subjects with autism spectrum disorders [15], which may be linked to the elevated testosterone effects on arousal and social anxiety [55], and contribute to male predominance in autism.

Why are there gender differences in brain development and function? Genetic information encoded in the sex chromosome has been suggested as an underlying reason. For example, the *SRY* gene on Y chromosome underlies sex-dependent neuroanatomical structures in various brain regions, as well as behavioral phenotypes, such as social exploration and aggression [21,25]. Other theories suggest that females' additional X-chromosome may provide the pattern of X-linked gene effects that is different from males [19].

Besides genomic influence, epigenetic factors also play a key role. Estradiol regulates its own receptor function via DNA methylation on the promoters of ERs, which is thought to contribute to the sexual dimorphism of neuronal anatomy and behavior [50]. Consistent with this, knockdown of ER with antisense oligonucleotides disrupts sexual differentiation of the brain [38]. ER knockout female mice display diminished maternal behaviors and exhibit male-like aggressive behaviors [51].

Postnatal stress is thought to affect the ability to cope with adversity in adulthood [3]. Emerging evidence suggests that the vulnerability to stress in developing brains is also gender-dependent. Study of prenatal stress shows that male, but not female, offspring exposed to stress in early pregnancy have maladaptive stress responses, which is associated with alterations in corticotrophin-releasing factor (CRF) and glucocorticoid receptor (GR) expression [47]. Female offspring is more resilient to various early life stress challenges than male offspring [35]. When exposed to postnatal stress, such as early weaning from maternal lactation or limited nesting & bedding materials, neurogenesis is perturbed in male, but not female, animals [29,49]. Such gender-dependent susceptibility to stress during development may be linked to mental illnesses in adult life [4,19].

In summary, animal and human studies have found that males and females show distinct stress responses and have different vulnerability to stress-related mental disorders. Recent studies




Fig. 2. A diagram showing the role of estrogen in determining the sexually dimorphic effects of stress on glutamatergic synaptic function in prefrontal cortical neurons. In males, repeated stress triggers the activation of glucocorticoid receptors (GR). GR bind to glucocorticoid response element (GRE) on the promoters of downstream genes, triggering the increased glutamate receptor ubiquitination and degradation. In females, estrogen activates estrogen receptors (ER), which bind to estrogen response element (ERE) and enhance the transcription of synaptic plasticity genes that promote the synthesis and exocytosis of glutamate receptors, therefore counteracting the stress-induced depressing effects.

have suggested the contribution of estrogen in the brain to this sexual dimorphism. The differential effects of stress on glutamatergic transmission in males vs. females, which are attributed to the influence of estrogen on synaptic plasticity (Fig. 2, [77,70], may underlie the sex-specific impact of stress on cognitive processes. More detailed genetic, epigenetic and molecular mechanisms await to be elucidated.

Acknowledgement

This work was supported by NIH grant (R01-DA037618) to Z.Y.

References

- [1] M.M. Adams, S.E. Fink, R.A. Shah, W.G. Janssen, S. Hayashi, T.A. Milner, B.S. McEwen, J.H. Morrison, Estrogen and aging affect the subcellular distribution of estrogen receptor- α in the hippocampus of female rats, *J. Neurosci.* 22 (2002) 3608–3614.
- [2] K.T. Akama, B.S. McEwen, Estrogen stimulates postsynaptic density-95 rapid protein synthesis via the Akt/protein kinase B pathway, *J. Neurosci.* 23 (2003) 2333–2339.
- [3] R.C. Bagot, F.N. van Hasselt, D.L. Champagne, M.J. Meaney, H.J. Krugers, M. Joels, Maternal care determines rapid effects of stress mediators on synaptic plasticity in adult rat hippocampal dentate gyrus, *Neurobiol. Learn. Mem.* 92 (2009) 292–300.
- [4] T.L. Bale, Neuroendocrine and immune influences on the CNS: it's a matter of sex, *Neuron* 64 (2009) 13–16.
- [5] D.A. Bangasser, A. Curtis, B.A. Reyes, T.T. Bethea, I. Parastatidis, H. Ischiropoulos, E.J. Van Bockstaele, R.J. Valentino, Sex differences in corticotropin-releasing factor receptor signaling and trafficking: potential role in female vulnerability to stress-related psychopathology, *Mol. Psychiatry* 15 (877) (2010) 896–904.
- [6] K.D. Beck, V.N. Luine, Food deprivation modulates chronic stress effects on object recognition in male rats: role of monoamines and amino acids, *Brain Res.* 830 (1999) 56–71.
- [7] K.D. Beck, V.N. Luine, Sex differences in behavioral and neurochemical profiles after chronic stress: role of housing conditions, *Physiol* 75 (2002) 661–673.
- [8] R.E. Bowman, Stress-induced changes in spatial memory are sexually differentiated and vary across the lifespan, *J. Neuroendocrinol.* 17 (2005) 526–535.
- [9] R.E. Bowman, K.D. Beck, V.N. Luine, Chronic stress effects on memory. Sex differences in performance and monoamines, *Horm. Behav.* 43 (2003) 48–59.
- [10] R.E. Bowman, D. Ferguson, V.N. Luine, Effects of chronic restraint stress and estradiol on open field activity, spatial memory, and monoaminergic neurotransmitters in ovariectomized rats, *Neuroscience* 113 (2002) 401–410.
- [11] R.E. Bowman, R. Micik, C. Gautreaux, L. Fernandez, V.N. Luine, Sex-dependent changes in anxiety, memory, and monoamines following one week of stress, *Physiol. Behav.* 97 (2009) 21–29.
- [12] R.E. Bowman, M.C. Zrull, V.N. Luine, Chronic restraint stress enhances radial arm maze performance in female rats, *Brain Res.* 904 (2001) 279–289.
- [13] T.M. Bredemann, L.L. McMahon, 17 β Estradiol increases resilience and improves hippocampal synaptic function in helpless ovariectomized rats, *Psychoneuroendocrinology* 42 (2014) 77–88.
- [14] N. Breslau, H.D. Chilcoat, R.C. Kessler, E.L. Peterson, V.C. Lucia, Vulnerability to assaultive violence: further specification of the sex difference in post-traumatic stress disorder, *Psychol. Med.* 29 (1999) 813–821.
- [15] A. Crider, R. Thakkar, A.O. Ahmed, A. Pillai, Dysregulation of estrogen receptor beta (ER β), aromatase (CYP19A1), and ER co-activators in the middle frontal gyrus of autism spectrum disorder subjects, *Mol. Autism.* 5 (2014) 46.
- [16] H. Cohen, R. Yehuda, Gender differences in animal models of posttraumatic stress disorder, *Dis. Markers* 30 (2011) 141–150.
- [17] C.D. Conrad, L.A. Galea, Y. Kuroda, B.S. McEwen, Chronic stress impairs rat spatial memory on the Y maze, and this effect is blocked by tianeptine pretreatment, *Behav. Neurosci.* 110 (1996) 1321–1334.
- [18] C.D. Conrad, K.D. Grote, R.J. Hobbs, A. Ferayorni, Sex differences in spatial and non-spatial Y-maze performance after chronic stress, *Neurobiol. Learn. Mem.* 79 (2003) 32–40.

- [19] E.P. Davis, D. Pfaff, Sexually dimorphic responses to early adversity: implications for affective problems and autism spectrum disorder, *Psychoneuroendocrinology* 49 (2014) 11–25.
- [20] E.R. de Kloet, M. Joëls, F. Holsboer, Stress and the brain: from adaptation to disease, *Nat. Rev. Neurosci.* 6 (2005) 463–475.
- [21] G.J. De Vries, E.F. Rissman, R.B. Simerly, L.Y. Yang, E.M. Scordalakes, C.J. Auger, A. Swain, R. Lovell-Badge, P.S. Burgoyne, A.P. Arnold, A model system for study of sex chromosome effects on sexually dimorphic neural and behavioral traits, *J. Neurosci.* 22 (2002) 9005–9014.
- [22] D.M. Diamond, M.C. Bennett, M. Fleshner, G.M. Rose, Inverted-U relationship between the level of peripheral corticosterone and the magnitude of hippocampal primed burst potentiation, *Hippocampus* 2 (1992) 421–430.
- [23] M.L. Gomez, L. Martinez-Mota, E. Estrada-Camarena, A. Fernandez-Guasti, Influence of the brain sexual differentiation process on despair and antidepressant-like effect of fluoxetine in the rat forced swim test, *Neuroscience* 261 (2014) 11–22.
- [24] T. Hajszan, K. Szigeti-Buck, N.L. Sallam, J. Bober, A. Parducz, N.J. Macluskus, C. Leranth, R.S. Duman, Effects of estradiol on learned helplessness and associated remodeling of hippocampal spine synapses in female rats, *Biol. Psychiatry* 67 (2010) 168–174.
- [25] R.A. Morsbroek, F. Sluyter, P.V. Guillot, G.A. Van Oortmerssen, W.E. Crusio, Y chromosomal effects on hippocampal mossy fiber distributions in mice selected for aggression, *Brain Res.* 682 (1995) 203–206.
- [26] Y. Hojo, T.-A. Hattori, T. Enami, A. Furukawa, K. Suzuki, et al., Adult male rat hippocampus synthesizes estradiol from pregnenolone by cytochromes P45017a and P450 aromatase localized in neurons, *Proc. Natl. Acad. Sci. USA* 101 (2003) 865–870.
- [27] M. Joels, Corticosteroid effects in the brain: U-shape it, *Trends Pharmacol. Sci.* 27 (2006) 244–250.
- [28] I.N. Karatsoreos, B.S. McEwen, Psychobiological allostasis: resistance, resilience and vulnerability, *Trends Cogn. Sci.* 15 (2011) 576–584.
- [29] T. Kikusui, Y. Mori, Behavioural and neurochemical consequences of early weaning in rodents, *J. Neuroendocrinol.* 21 (2009) 427–431.
- [30] T. Kimoto, T. Tsurugizawa, Y. Ohta, J. Makino, H. Tamura, Y. Hojo, N. Takata, S. Kawato, Neurosteroid synthesis by cytochrome p450-containing systems localized in the rat brain hippocampal neurons: N-methyl-D-aspartate and calcium-dependent synthesis, *Endocrinology* 142 (2001) 3578–3589.
- [31] E. Kitraki, O. Kremmyda, D. Youlatos, M.N. Alexis, C. Kittas, Gender-dependent alterations in corticosteroid receptor status and spatial performance following 21 days of restraint stress, *Neuroscience* 125 (2004) 47–55.
- [32] A.T. Konkle, M.M. McCarthy, Developmental time course of estradiol, testosterone, and dihydrotestosterone levels in discrete regions of male and female rat brain, *Endocrinology* 152 (2011) 223–235.
- [33] E.A. Kramár, L.Y. Chen, N.J. Brandon, C.S. Rex, F. Liu, C.M. Gall, G. Lynch, Cytoskeletal changes underlie estrogen's acute effects on synaptic transmission and plasticity, *J. Neurosci.* 29 (2009) 12982–12993.
- [34] G.S. Kranz, A. Hahn, U. Kaufmann, M. Kublbock, A. Hummer, S. Ganger, R. Seiger, D. Winkler, D.F. Swaab, C. Windischberger, et al., White matter microstructure in transsexuals and controls investigated by diffusion tensor imaging, *J. Neurosci.* 34 (2014) 15466–15475.
- [35] N. Lajud, L. Torner, Early life stress and hippocampal neurogenesis in the neonate: sexual dimorphism, long term consequences and possible mediators, *Front. Mol. Neurosci.* 8 (2015) 3.
- [36] Y.L. Lan, J. Zhao, S. Li, Update on the neuroprotective effect of estrogen receptor alpha against Alzheimer's disease, *J. Alzheimers Dis.* 43 (2015) 1137–1148.
- [37] V.N. Luine, K.D. Beck, R.E. Bowman, M. Frankfurt, N.J. Macluskus, Chronic stress and neural function: accounting for sex and age, *J. Neuroendocrinol.* 19 (2007) 743–751.
- [38] M.M. McCarthy, E.H. Schlenker, D.W. Pfaff, Enduring consequences of neonatal treatment with antisense oligodeoxynucleotides to estrogen receptor messenger ribonucleic acid on sexual differentiation of rat brain, *Endocrinology* 133 (1993) 433–439.
- [39] L.D. McCullough, K. Blizard, E.R. Simpson, O.K. Oz, P.D. Hurn, Aromatase cytochrome P450 and extragonadal estrogen play a role in ischemic neuroprotection, *J. Neurosci.* 23 (2003) 8701–8705.
- [40] B.S. McEwen, Stress, sex, and neural adaptation to a changing environment: mechanisms of neuronal remodeling, *Ann. N. Y. Acad. Sci.* 1204 (Suppl) (2010) 38–59.
- [41] B.S. McEwen, K.T. Akama, J.L. Spencer-Segal, T.A. Milner, E.M. Waters, Estrogen effects on the brain: actions beyond the hypothalamus via novel mechanisms, *Behav. Neurosci.* 126 (2012) 4–16.
- [42] B. McEwen, K. Akama, S. Alves, W.G. Brake, K. Bulloch, S. Lee, C. Li, G. Yuen, T.A. Milner, Tracking the estrogen receptor in neurons: implications for estrogen-induced synapse formation, *Proc. Natl. Acad. Sci. USA* 98 (2001) 7093–7100.
- [43] B.S. McEwen, Physiology and neurobiology of stress and adaptation: central role of the brain, *Physiol. Rev.* 87 (2007) 873–904.
- [44] K.F. McLaughlin, S.E. Baran, R.L. Wright, C.D. Conrad, Chronic stress enhances spatial memory in ovariectomized female rats despite CA3 dendritic retraction. Possible involvement of CA1 neurons, *Neuroscience* 135 (2005) 1045–1054.
- [45] M.R. Milad, S.A. Igoe, K. Lebron-Milad, J.E. Novales, Estrous cycle phase and gonadal hormones influence conditioned fear extinction, *Neuroscience* 164 (2009) 887–895.
- [46] M.R. Milad, M.A. Zeidan, A. Contero, R.K. Pitman, A. Klibanski, S.L. Rauch, et al., The influence of gonadal hormones on conditioned fear extinction in healthy humans, *Neuroscience* 168 (2010) 652–658.
- [47] B.R. Mueller, T.L. Bale, Sex-specific programming of offspring emotionality after stress early in pregnancy, *J. Neurosci.* 28 (2008) 9055–9065.
- [48] D.D. Murphy, N.B. Cole, M. Segal, Brain-derived neurotrophic factor mediates estradiol-induced dendritic spine formation in hippocampal neurons, *Proc. Natl. Acad. Sci. USA* 95 (1998) 11412–11417.
- [49] E.F. Naninck, L. Hoeijmakers, N. Kakava-Georgiadou, A. Meesters, S.E. Lasic, P.J. Lucassen, A. Korosi, Chronic early life stress alters developmental and adult neurogenesis and impairs cognitive function in mice, *Hippocampus* 25 (2015) 309–328.
- [50] B.M. Nugent, J.M. Schwarz, M.M. McCarthy, Hormonally mediated epigenetic changes to steroid receptors in the developing brain: implications for sexual differentiation, *Horm. Behav.* 59 (2011) 338–344.
- [51] S. Ogawa, J.A. Taylor, D.B. Lubahn, K.S. Korach, D.W. Pfaff, Reversal of sex roles in genetic female mice by disruption of estrogen receptor gene, *Neuroendocrinology* 64 (1996) 467–470.
- [52] B.K. Ormerod, T.T. Lee, L.A. Galea, Estradiol enhances neurogenesis in the dentate gyri of adult male meadow voles by increasing the survival of young granule neurons, *Neuroscience* 128 (2004) 645–654.
- [53] S.E. Pérez, E.Y. Chen, E.J. Mufson, Distribution of estrogen receptor alpha and beta immunoreactive profiles in the postnatal rat brain, *Brain Res. Dev. Brain Res.* 145 (2003) 117–139.
- [54] M.J. Perez-Alvarez, L. Mateos, A. Alonso, F. Wandosell, Estradiol and Progesterone Administration After pMCAO Stimulates the Neurological Recovery and Reduces the Detrimental Effect of Ischemia Mainly in Hippocampus, *Mol. Neurobiol.* (2014) [epub ahead of print].
- [55] D.W. Pfaff, I. Rapin, S. Goldman, Male predominance in autism: neuroendocrine influences on arousal and social anxiety, *Autism Res.* 4 (2011) 163–176.
- [56] F.H. Pottou, M. Bhowmik, D. Vohora, Raloxifene protects against seizures and neurodegeneration in a mouse model mimicking epilepsy in postmenopausal woman, *Eur. J. Pharm. Sci.* 65C (2014) 167–173.
- [57] P.V. Rekkas, A.A. Wilson, V.W. Lee, P. Yagalasingam, J. Sacher, P. Rusjan, S. Houle, D.E. Stewart, N.J. Kolla, S. Kish, L. Chiuccariello, J.H. Meyer, Greater monoamine oxidase a binding in perimenopausal age as measured with carbon 11-labeled harmine positron emission tomography, *JAMA Psychiatry* 71 (2014) 873–879.
- [58] C.D. Rey, J. Lipps, R.M. Shansky, Dopamine D1 receptor activation rescues extinction impairments in low-estrogen female rats and induces cortical layer-specific activation changes in prefrontal-amygdala circuits, *Neuropsychopharmacology* 39 (2014) 1282–1289.
- [59] K. Sellers, P. Raval, D.P. Srivastava, Molecular signature of rapid estrogen regulation of synaptic connectivity and cognition, *Front. Neuroendocrinol.* (2014) [epub ahead of print].
- [60] R.M. Shansky, C. Glavis-Bloom, D. Lerman, P. McRae, C. Benson, K. Miller, L. Cosand, T.L. Horvath, A.F. Arnsten, Estrogen mediates sex differences in stress-induced prefrontal cortex dysfunction, *Mol. Psychiatry* 9 (2004) 531–538.
- [61] R.M. Shansky, C. Hamo, P.R. Hof, W. Lou, B.S. McEwen, J.H. Morrison, Estrogen promotes stress sensitivity in a prefrontal cortex-amygdala pathway, *Cereb. Cortex* 20 (2010) 2560–2567.
- [62] T.J. Shors, C. Chua, J. Falduto, Sex differences and opposite effects of stress on dendritic spine density in the male versus female hippocampus, *J. Neurosci.* 21 (2001) 6292–6297.
- [63] T. Smejkalova, C.S. Woolley, Estradiol acutely potentiates hippocampal excitatory synaptic transmission through a presynaptic mechanism, *J. Neurosci.* 30 (2010) 16137–16148.
- [64] J.L. Spencer, E.M. Waters, R.D. Romeo, G.E. Wood, T.A. Milner, B.S. McEwen, Uncovering the mechanisms of estrogen effects on hippocampal function, *Front. Neuroendocrinol.* 29 (2008) 219–237.
- [65] L. Sterrenburg, B. Gaszner, J. Boerrigter, L. Santbergen, M. Bramini, E. Elliott, A. Chen, B.W. Peeters, E.W. Roubos, T. Kozicz, Chronic stress induces sex-specific alterations in methylation and expression of corticotropin-releasing factor gene in the rat, *PLoS One* 6 (2011) e28128.
- [66] S. Suda, E. Segi-Nishida, S.S. Newton, R.S. Duman, A postpartum model in rat: behavioral and gene expression changes induced by ovarian steroid deprivation, *Biol. Psychiatry* 64 (2008) 311–319.
- [67] V. Vialou, J. Feng, A.J. Robison, E.J. Nestler, Epigenetic mechanisms of depression and antidepressant action, *Annu. Rev. Pharmacol. Toxicol.* 53 (2013) 59–87.
- [68] A.A. Wolf, C.A. Frye, A review and update of mechanisms of estrogen in the hippocampus and amygdala for anxiety and depression behavior, *Neuropsychopharmacology* 31 (2006) 1097–1111.
- [69] A.C. Wang, Y. Hara, W.G. Janssen, P.R. Rapp, J.H. Morrison, Synaptic estrogen receptor-alpha levels in prefrontal cortex in female rhesus monkeys and their correlation with cognitive performance, *J. Neurosci.* 30 (2010) 12770–12776.
- [70] J. Wei, E.Y. Yuen, W. Liu, X. Li, P. Zhong, I.N. Karatsoreos, B.S. McEwen, Z. Yan, Estrogen protects against the detrimental effects of repeated stress on glutamatergic transmission and cognition, *Mol. Psychiatry* 19 (2014) 588–598.
- [71] M.M. Weissman, R.C. Bland, G.J. Canino, C. Faravelli, S. Greenwald, H.G. Hwu, P. R. Joyce, E.G. Karam, C.K. Lee, J. Lellouch, et al., Cross-national epidemiology of major depression and bipolar disorder, *JAMA* 276 (1996) 293–299.
- [72] C.S. Woolley, Acute effects of estrogen on neuronal physiology, *Annu. Rev. Pharmacol. Toxicol.* 47 (2007) 657–680.
- [73] C.S. Woolley, N.G. Weiland, B.S. McEwen, P.A. Schwartzkroin, Estradiol increases the sensitivity of hippocampal CA1 pyramidal cells to NMDA receptor-mediated synaptic input: correlation with dendritic spine density, *J. Neurosci.* 17 (1997) 1848–1859.

- [74] H. Yu, D.D. Wang, Y. Wang, T. Liu, F.S. Lee, Z.Y. Chen, Variant brain-derived neurotrophic factor Val66Met polymorphism alters vulnerability to stress and response to antidepressants, *J. Neurosci.* 32 (2012) 4092–4101.
- [75] E.Y. Yuen, W. Liu, I.N. Karatsoreos, J. Feng, B.S. McEwen, Z. Yan, Acute stress enhances glutamatergic transmission in prefrontal cortex and facilitates working memory, *Proc. Natl. Acad. Sci. USA* 106 (2009) 14075–14079.
- [76] E.Y. Yuen, W. Liu, I.N. Karatsoreos, Y. Ren, J. Feng, B.S. McEwen, Z. Yan, Mechanisms for acute stress-induced enhancement of glutamatergic transmission and working memory, *Mol. Psychiatry* 16 (2011) 156–170.
- [77] E.Y. Yuen, J. Wei, W. Liu, P. Zhong, X. Li, Z. Yan, Repeated stress causes cognitive impairment by suppressing glutamate receptor expression and function in prefrontal cortex, *Neuron* 73 (2012) 962–977.
- [78] J.M. Zhang, A.T. Konkle, S.L. Zup, M.M. McCarthy, Impact of sex and hormones on new cells in the developing rat hippocampus: a novel source of sex dimorphism? *Eur. J. Neurosci.* 27 (2008) 791–800.